



AIM: AI数学家系统

李鹏

清华大学

AI深刻改变科学研究范式



清华大学
Tsinghua University


- 深度学习的发展赋能AI4Science，极大地推动了科学的发展。



John J. Hopfield and Geoffrey Hinton. Ill. Niklas Elmehed © Nobel Prize Outreach




David Baker, Demis Hassabis and John Jumper. Ill. Niklas Elmehed © Nobel Prize Outreach




AlphaQuant

Solution
Construct D: midpoint BH [a]
[a], O_2 midpoint HQ $\Rightarrow$ BQ $\parallel$ O_2D [20]
...
Construct G: midpoint HC [b] ...
 $\angle GMD = \angle GO_2D \Rightarrow M O_2 G D$ cyclic [26]
[a], [b] $\Rightarrow$ BC $\parallel$ DG [30]
...
Construct E: midpoint MK [c]
..., [c] $\Rightarrow \angle KFC = \angle KO_1E$ [104]
...
 $\angle FK O_1 = \angle FK O_2 \Rightarrow K O_1 \parallel K O_2$ [109]
[109] $\Rightarrow O_1 O_2 K$ collinear $\Rightarrow (O_1)(O_2)$ tangent



AlphaGeometry



AlphaFold


<https://www.nobelprize.org/all-nobel-prizes-2024/>
<https://www.nature.com/articles/s41586-024-07487-w>
<https://www.nature.com/articles/s41586-024-08148-8>
<https://www.nature.com/articles/s41586-023-06747-5>

数学研究意义重大




清华大学
Tsinghua University


- 数学研究具有重要的理论与应用价值，是人类智慧的集中体现



AI与数学的结合可以产生什么样的化学反应？



探索至真至美的自然真理





科学研究与工程应用的理论工具

数学之于AI：大模型性能进步关键要素



清华大学
Tsinghua University

● 数学和代码是当前提升大模型推理能力的关键要素



针对数学和代码任务训练的第一个推理模型
o1能够在各方面取得明显进步

DeepSeek-R1通过针对数学问题的可验证反馈的强化学习训练获取了深度思考的能力

AI之于数学：数学能力快速接近人类



清华大学
Tsinghua University

● 2025年高考数学新课标 I 卷大模型最高水平达到145分（满分150分）

7家大模型挑战高考数学新一卷				
模型	客观题（文本）得分	图像题（单选）得分	解答题得分	总分
Gemini 2.5 pro	68	0	77	145
Doubao-1.5-thinking-vision-pro	68	0	76	144
DeepSeek R1	68	/	76	144
o3	65	0	75	140
Qwen3-235b	68	/	71	139
hunyuan-t1-latest	68	/	68	136
文心X1 Turbo	68	/	66	134
注：本次评测分为客观题（文本）、第6题图像题（单选）和解答题，客观题（文本）总计68分，图像题总共（单选）5分，解答题77分，总分150分				


图像理解能力不足导致失分

AI之于数学：数学能力快速接近人类



清华大学
Tsinghua University

● 2025年5月o4-mini在部分前沿数学问题上的表现已经达到专家水平




“我不想加剧恐慌。但在某些方面，大语言模型的表现已经超越了世界上大多数最优秀的研究生。”

——小野健

o4-mini outperforms the average team of mathematicians on our FrontierMath human baseline competition

EPOCH AI



Competition problems are taken from a non-random subset of FrontierMath and tailored to the background knowledge of contestants. Participants were grouped into 8 human teams with 4-5 mathematicians each. The "All teams" score is determined by looking at how many problems were correctly answered by at least one team, though note that each team only attempted around 10 out of 23 problems on average.

CC-BY

epoch.ai

AIME: 高中竞赛水平测试集



清华大学
Tsinghua University

- AIME数据集来自于美国数学邀请赛，赛事面向高中生，是选拔国际数学奥林匹克美国队的赛事之一

赛题设置

- 共15道题，每一题的难度不断增大
- 答案均为一个介于0-999之间的整数

局限性

- 答案存在且已知
- 主要考查解题技巧

Problem

The 9 members of a baseball team went to an ice-cream parlor after their game. Each player had a single scoop cone of chocolate, vanilla, or strawberry ice cream. At least one player chose each flavor, and the number of players who chose chocolate was greater than the number of players who chose vanilla, which was greater than the number of players who chose strawberry. Let N be the number of different assignments of flavors to players that meet these conditions. Find the remainder when N is divided by 1000.

Solution 1

Let c be the number of players who choose chocolate, v be the number of players who choose vanilla, and s be the number of players who choose strawberry ice cream. We are given two pieces of information $c, v, s \geq 1$ and $c + v + s = 9$.

By inspection the only solutions for (c, v, s) are $(2, 3, 4)$, $(1, 2, 6)$, $(1, 3, 5)$.

Now we must choose which player chooses which flavor. For the general case (c, v, s) , we begin by choose c of the 9 players who eat chocolate, then we choose v of the $9 - c$ players who vanilla, after this the amount of players who eat strawberry is fixed. Therefore the general

formula is $\binom{9}{c} \binom{9-c}{v}$.

Therefore our final answer is,

$$\binom{9}{2} \binom{7}{3} + \binom{9}{1} \binom{8}{2} + \binom{9}{1} \binom{8}{3} = 2\boxed{016}.$$

AIME: 高中竞赛水平测试集



清华大学
Tsinghua University

- 当前大模型主要使用可验证的练习题或者数学竞赛题用于训练和测试，与AIME题目类型分布一致性强

DeepSeek-R1针对AIME2025竞赛的第一题给出的完整正确解答

- **原始题目:** Find the sum of all integer bases $b > 9$ for which 17_b is a divisor of 97_b
- **题目难度:** 结构简单且不涉及高深概念
- **思考长度:** 可在简短步骤当中快速完成
- **考查方面:** 主要考查解题技解的应用



The problem requires finding all integer bases $b > 9$ such that 17_b divides 97_b , and then summing those bases.

Convert 17_b and 97_b to base 10:

- $17_b = 1 \cdot b + 7 = b + 7$
- $97_b = 9 \cdot b + 7$

The condition that 17_b divides 97_b means that $b + 7$ divides $9b + 7$, or $(b + 7) \mid (9b + 7)$.

Set $d = b + 7$, so $b = d - 7$. Substitute into $9b + 7$:

$$9b + 7 = 9(d - 7) + 7 = 9d - 63 + 7 = 9d - 56$$

The condition becomes $d \mid (9d - 56)$. Since $d \mid 9d$, it follows that $d \mid (9d - (9d - 56)) = d \mid 56$.

Given $b > 9$, it follows that $d = b + 7 > 16$. Thus, d must be a positive divisor of 56 greater than 16.

The positive divisors of 56 are 1, 2, 4, 7, 8, 14, 28, and 56. The divisors greater than 16 are 28 and 56.

- If $d = 28$, then $b = 28 - 7 = 21$.
- If $d = 56$, then $b = 56 - 7 = 49$.

Both bases satisfy $b > 9$. Verify the division condition:

- For $b = 21$:
 - $17_{21} = 1 \cdot 21 + 7 = 28$
 - $97_{21} = 9 \cdot 21 + 7 = 189 + 7 = 196$
 - $196 \div 28 = 7$, which is an integer.
- For $b = 49$:
 - $17_{49} = 1 \cdot 49 + 7 = 56$
 - $97_{49} = 9 \cdot 49 + 7 = 441 + 7 = 448$
 - $448 \div 56 = 8$, which is an integer.

Both bases are valid, and there are no other bases $b > 9$ that satisfy the condition. The digits in both numbers are valid since $b > 9$ implies $b > 9$ for digit 9 in 97_b and $b > 7$ for digit 7 in both numbers.

The sum of the bases is $21 + 49 = 70$.

`\boxed{70}`

FrontierMath: 挑战性陡升



清华大学
Tsinghua University

- 由专业数学家撰写，陶哲轩等均认可FrontierMath具有高挑战性，但仍与研究级数学问题存在显著差距

挑战度高

- 题目由专业数学家撰写
- 涵盖现代数学主要分支
- 人类专家一般需要数小时解决一个题目

局限性

- 答案存在且已知
- 难度低于数学研究问题

Construct a degree 19 polynomial $p(x) \in \mathbb{C}[x]$ such that $X := \{p(x) = p(y)\} \subset \mathbb{P}^1 \times \mathbb{P}^1$ has at least 3 (but not all linear) irreducible components over $\mathbb{C}$. Choose $p(x)$ to be odd, monic, have real coefficients and linear coefficient -19 and calculate $p(19)$.

Answer: 1876572071974094803391179

MSC classification: 14 Algebraic geometry; 20 Group theory and generalizations; 11 Number theory generalizations

数学解题 v.s. 数学研究



差异1：长线思考

数学研究需要比做题显著更多、更加长期且更加深入的有效思考

三维挂谷猜想

- 完整证明总计12个章节、125页正文
- 论文仅呈现了最终证明过程，真实思考和推理量远大于论文呈现结果

Volume estimates for unions of convex sets, and the Kakeya set conjecture in three dimensions	
Hong Wang* Joshua Zahl †	
February 26, 2025	
Abstract	
We study sets of δ tubes in $\mathbb{R}^3$, with the property that not too many inside a common convex set V . We show that the union of tubes fro almost maximal volume. As a consequence, we prove that every Kakeya and Hausdorff dimension 3.	
Contents	
1 Introduction	23
1.1 Theorem 1.2 and multi-scale analysis	21
1.2 Unions of convex sets, and non-clustering	22
1.3 From Assertions $\mathcal{D}$ and $\mathcal{E}$ to the Kakeya set conjecture	23
1.4 Proof philosophy, and previous work on the Kakeya set conjecture	23
1.5 A vignette of the proof	23
1.6 Tube doubling and Keleti's line segment extension conjecture	24
1.7 Thanks	24
2 A sketch of the proof	23
2.1 Proposition 1.6: Assertions $\mathcal{D}$ and $\mathcal{E}$ are equivalent	23
2.2 A two-scale grains decomposition	23
2.3 Refined induction on scales	23
2.4 Multi-scale structure, Nikishin-Stein-Pisier factorization, and Sticky Kakeya	24
3 Notation	23
3.1 Convex sets and shadings	23
3.2 Table of notation	24
4 Wolff Axioms and Factoring Convex Sets	25
4.1 Definitions: Wolff axioms and covers	25
4.2 Factoring Convex Sets	26
4.3 Convex Sets and the Frostman Slab Wolff Axioms	31
4.4 The Frostman Slab Wolff Axioms and Covers	31
5 Factoring tubes into flat prisms	25
5.1 A few frequently used Cordoba-type L^2 arguments	25
5.1.1 A volume estimate for slabs	25
5.1.2 Tangential vs transverse prism intersection	26
5.2 Assertions $\mathcal{F}$, $\mathcal{E}$, and $\tilde{\mathcal{E}}$ are equivalent	26
5.3 Proof of Proposition 5.1: Tubes that factor through flat boxes	26
5.4 Proof of Proposition 5.2: Factoring at two scales	26
5.5 Tubes organized into to slabs	26
6 Assertions $\mathcal{D}$ and $\mathcal{E}$ are equivalent	26
6.1 Proof of Proposition 6.3: A factoring trichotomy	26
7 A two-scale grains decomposition for tubes in $\mathbb{R}^3$	60
7.1 Broadness	63
7.2 Broadness and the Frostman Slab Wolff axioms	66
7.3 The iteration base case: Guth's grains decomposition	69
7.4 Moves #1, #2, #3: Parallel structure	70
7.5 Using Moves #1, #2, #3 to prove Proposition 7.5	72
8 Moves #1, #2, and #3	74
8.1 Move #1: Replacing grains with longer grains to ensure $c \geq \delta_{\delta}^{\mathcal{E}}(\#T_{\rho})/(\#T)$	74
8.2 Move #2: Replacing square grains with longer grains	74
8.3 Move #3: Replacing grains with wider grains with small C_{KT-CW}^{loc}	85
9 A refined induction-on-scales argument	100
10 Sticky Kakeya for tubes satisfying the Katz-Tao Convex Wolff Axioms at every Scale	103
10.1 Nikishin-Stein-Pisier Factorization and the Convex Wolff Axioms	105
11 Multi-scale analysis and the proof of Proposition 1.7	110
12 Tube Doubling	114

*Courant Institute of Mathematical Sciences, New York University, New York, NY, USA.
†Department of Mathematics, The University of British Columbia, Vancouver, BC, Canada.

数学解题 v.s. 数学研究



差异2：过程严谨

错误零容忍，正确性需要可验证，而验证代价异常高昂

Hence if $\mathcal{W}' \subset \mathcal{W}_0$, to compare $\# \left(\bigcup_{W_i \in \mathcal{W}'} \mathcal{U}_0[W_i] \right)$ and $\sum_{W_i \in \mathcal{W}'} \# \mathcal{U}_0[W_i]$,

三维挂谷猜想
证明片段

$$\begin{aligned} \kappa_0 \frac{C_{KT-CW}(\mathcal{U}_0)}{|\mathcal{U}|} \sum_{W_i \in \mathcal{W}'} |W_i| &\leq \sum_{W_i \in \mathcal{W}'} \# \mathcal{U}_{i-1}[W_i] = \# \left(\bigsqcup_{W_i \in \mathcal{W}'} \mathcal{U}_{i-1}[W_i] \right) \\ &\leq \# \left(\bigcup_{W_i \in \mathcal{W}'} \mathcal{U}_0[W_i] \right) \leq \sum_{W_i \in \mathcal{W}'} \# \mathcal{U}_0[W_i] \leq \frac{C_{KT-CW}(\mathcal{U}_0)}{|\mathcal{U}|} \sum_{W_i \in \mathcal{W}'} |W_i|. \end{aligned} \tag{4.12}$$

The equality in (4.12) uses the critical fact that if $i \neq i'$, then $\mathcal{U}_{i-1}[W_i]$ and $\mathcal{U}_{i'-1}[W_{i'}]$ are disjoint.

案例	验证过程	验证耗时
费马大定理	1993年给出初始证明，由一个顶级小组反复审读并指出问题，随后怀尔斯与他的学生泰勒合作修改，1995年才正式发表两篇论文	约2年
庞加莱猜想	2002-2003年分三篇预印本形式在线发布了证明，数学界以非正式审稿方式组织了由多位几何拓扑学家验证、重写和解释其证明的过程	约4-5年

数学解题 v.s. 数学研究




清华大学
Tsinghua University

差异3：答案存在性

数学解题答案存在性**已知**且**确定**

数学研究答案存在性**未知**且**不确定**



哥德尔

- **第一不完备定理**：任何自治的形式系统，只要蕴涵皮亚诺算术公理，就可以在其中构造在体系中不能被证明的真命题，因此通过推理演绎不能得到所有真命题（即体系是不完备的）
- **第二不完备定理**：任何逻辑自治的形式系统，只要蕴涵皮亚诺算术公理，它就不能用于证明其本身的自治性

AI + 数学研究：形式化证明



清华大学
Tsinghua University

- 形式化证明历史悠久，国内外研究机构以及DeepSeek、字节跳动等积极探索该研究方向

优势

- 将各种数学证明转换为符号计算，可利用计算机机械化、批量执行
- 保证证明过程严格正确性

缺点

- 当前形式化证明工具存在明显的性能和效率问题
- 进而难以应用到前沿研究工作当中



Show that for any integer $n \geq 4$, we have $n^2 \leq n!$.

```
theorem induction_ineq_nsqlefactn (n : ℕ) (h₀ : 4 ≤ n) : n ^ 2 ≤ n ! := sorry
```

Understanding the Problem

We need to show that for any integer $n \geq 4$, the inequality $n^2 \leq n!$ holds. Here, $n!$ denotes the factorial of n , which is the product of all positive integers up to n .

Proof Sketch

To prove this formally, we can use mathematical induction. Here's how the induction would work:

1. **Base Case:** Verify the inequality for $n = 4$.
2. **Inductive Step:** Assume the inequality holds for some $k = 4$, i.e., $k^2 \leq k!$. Then show that it holds for $k + 1$, i.e., $(k + 1)^2 \leq (k + 1)!$.

chain-of-thought by DeepSeek-V3

Lean 4 Proof Structure

```
theorem induction_ineq_nsqlefactn (n : ℕ) (h₀ : 4 ≤ n) : n ^ 2 ≤ n ! := by
  have base_case : 4 ^ 2 ≤ 4 ! := by
    sorry
  have inductive_step : ∀ k ≥ 4, k ^ 2 ≤ k ! → (k + 1) ^ 2 ≤ (k + 1) ! := by
    sorry
  have final_proof : ∀ n ≥ 4, n ^ 2 ≤ n ! := by
    sorry
```



Recursively Solve Subgoals by DeepSeek-Prover-V2-7B

```
lemma inductive_step (n : ℕ) (h₀ : 4 ≤ n) (base_case : 4 ^ 2 ≤ 4 ! )
  : ∀ k ≥ 4, k ^ 2 ≤ k ! → (k + 1) ^ 2 ≤ (k + 1) ! := by
  sorry
```


Synthesize into Complete Formal Proofs

```
theorem induction_ineq_nsqlefactn (n : ℕ) (h₀ : 4 ≤ n) : n ^ 2 ≤ n ! := by
  have base_case : 4 ^ 2 ≤ 4 ! := by
    simp [Nat.factorial]
  have inductive_step : ∀ k ≥ 4, k ^ 2 ≤ k ! → (k + 1) ^ 2 ≤ (k + 1) ! := by
    intro k h₁ h₂
    simp_all [Nat.factorial]
    nlinarith
  have final_proof : ∀ n ≥ 4, n ^ 2 ≤ n ! := by
    intro n hn
    induction' hn with k hk
    case refl => exact base_case
    case step =>
      apply inductive_step k hk
      exact by assumption
  apply final_proof
  exact h₀
```

● 近期人们也开始探索利用大模型的能力执行数学科研任务的可能

AlphaEvolve

- **起源**：2025年5月14日由谷歌DeepMind推出
- **本质**：专注于算法优化的大模型智能体系统
- **成效**：已经能够独立取得一些新的研究成果
- **局限**：只适用于能够被转换为代码验证的问题



AIM: 面向数学研究的AI数学家智能体



清华大学
Tsinghua University

● AIM在利用人工智能进行复杂数学理论研究中迈出了重要的一步

数学解题

- 推理链条短
- 严谨性易满足
- 答案存在性已知
- 答案确定

数学研究

- 推理链条长
- 严谨性难满足
- 答案存在性未知
- 答案不确定

数学问题

量子算法问题

(结论已知)

Quantum Algorithm Problem

吸收边界问题

(结论已知)

Absorbing Boundary Condition

高对比度极限

(结论已知)

High Contrast Limit

均匀化问题

(结论未知)

Homogenization Problem

AIM效果

AIM完成了带有详细分析
推导过程的证明

AIM基本完成此问题的证
明过程和细节

AIM完成了主要结论的证
明过程，并且证明了新的
结论

AIM给出了合理的证明思
路和部分结论的证明，对
数学工作者很有启发意义

部分证明步骤

Multiply through by $e^{i\pi/4 - \epsilon t}$, cancel terms, and simplify:
$$\frac{d}{dt} \langle \psi, \psi \rangle + \frac{d}{dt} \langle \psi, \psi \rangle = 0 \implies \langle \psi, \psi \rangle = \langle \psi, \psi \rangle_0$$

Step 2: Spatial Discretization**
Discretize ψ on a grid with spacing Δx . Approximate $\partial_x^2 \psi$ via finite differences:
$$\partial_x^2 \psi \approx \frac{\psi(x+\Delta x) - 2\psi(x) + \psi(x-\Delta x))}{\Delta x^2}$$

Let D_h be the discretized second derivative matrix. The operator becomes:
$$D_h = -D_h$$

Step 3: Positive Semi-Definite Operator**
The matrix $D_h = -D_h$ is positive semi-definite because D_h is centered second derivative discretization is negative semi-definite. This aligns with the dispersive nature of the heat equation.
Step 4: Implications for Quantum Simulation**
While the original decomposition $D = \tilde{D} + \tilde{D}$ (with $\tilde{D} = -\tilde{D}$) is useful due to the absence of a livable term, the operator $\tilde{D}$ itself is suitable for quantum simulation methods designed for parabolic equations. Specifically, the time evolution $e^{-i\tilde{D}t}$ can be simulated using Trotterization or other discrete quantum algorithms, bypassing the need for the LDB terms.
[Correct] Variable substitution and equation transformation are correct. Discretization of the spatial variable is correct. And AIM finds the relevant PDE operators for the LDB terms. It's right!

2. The integral representation in the LDB terms for the BSM model can be approximated with precision using $O(1/\epsilon^2)$ terms through an adaptive discretization of the δ -integral, leveraging the rapid decay of the Cauchy kernel $\phi(\delta) = \frac{1}{\delta^2}$.
Proof: **Truncated Error Analysis** The integral $I = \int_{-\infty}^{\infty} \phi(\delta) \psi(x, \delta) d\delta$ is truncated to $[-R, R]$. The tail error is bounded by:
$$\left| \int_{| \delta | > R} \phi(\delta) \psi(x, \delta) d\delta \right| \leq \int_{| \delta | > R} \frac{1}{\delta^2} |\psi(x, \delta)| d\delta \leq \frac{1}{R} \max_{| \delta | > R} |\psi(x, \delta)|$$

Since $|\psi(x, \delta)| \leq C e^{-\delta^2/4}$, the tail error is $O(1/R)$. Thus, $R = O(1/\epsilon)$.
Adaptive Discretization The truncated integral $I_R = \int_{-R}^R \phi(\delta) \psi(x, \delta) d\delta$ is approximated using the trapezoidal rule with N nodes. The nodes are chosen as $\delta_k = \sqrt{4 \ln(2N)} e^{i\pi k/N}$. Since $\phi(\delta)$ and $\psi(x, \delta)$ are smooth, the truncation error is $O(1/N^2)$. The total error is $O(1/N^2) + O(1/N) = O(1/N)$.
Error Bound The truncated integral I_R includes all significant contributions, as the tail error is $O(1/N)$. The error bound is $O(1/N)$. The error bound is $O(1/N)$.
Time Step The time step Δt is chosen as $\Delta t = O(1/N^2)$. The error bound is $O(1/N^2)$.
Final Answer The numerical solution $\psi(x, t)$ is accurate to $O(1/N)$. The error bound is $O(1/N)$.
[Correct] Truncate and discretize the integral correctly.

Revised Proof of the a priori estimate Define the total energy functional:
$$E(t) = \frac{1}{2} \|\psi(t)\|_{L^2(\Omega)}^2 + \frac{1}{2} \|\psi(t)\|_{L^2(\Omega)}^2 = \frac{1}{2} \|\psi(t)\|_{L^2(\Omega)}^2$$

[Correct] The energy method and the correct energy functional form.

Step 1: Construct H^1 -Conforming Subspace Let $\{ \phi_j \}_{j=1}^N \subset H^1(\Omega)$ and $\{ \psi_j \}_{j=1}^N \subset H^1(\Omega)$ be eigenfunctions of the Laplace-Beltrami operators $-\Delta_{\Omega}$ and $-\Delta_{\Omega'}$ with Neumann/Dirichlet boundary conditions (as appropriate). These form an orthonormal basis for $L^2(\Omega)$ and $L^2(\Omega')$, respectively, and are dense in $H^1(\Omega)$ and $H^1(\Omega')$.
[Correct] The eigenfunctions form a basis for the function space.

**Step 2: Concretized Test with $w = \nabla \cdot \psi$ ** Substituting w into the variational equation:
$$\int_{\Omega} \nabla \cdot \psi \nabla \cdot \psi = \int_{\Omega} \nabla \cdot (\psi \nabla \cdot \psi) = \int_{\partial \Omega} \psi \nabla \cdot \psi \cdot \nu = 0$$

Step 3: Concretized Test with $w = \nabla \cdot \psi$ Substituting w into the variational equation:
$$\int_{\Omega} \nabla \cdot \psi \nabla \cdot \psi = \int_{\Omega} \nabla \cdot (\psi \nabla \cdot \psi) = \int_{\partial \Omega} \psi \nabla \cdot \psi \cdot \nu = 0$$

Rearranging and taking absolute values:
$$\left| \int_{\Omega} \nabla \cdot \psi \nabla \cdot \psi \right| \leq \int_{\Omega} |\nabla \cdot \psi|^2 = \int_{\Omega} |\nabla \cdot \psi|^2$$

Using Hölder inequality and trace dual:
$$\left| \int_{\Omega} \nabla \cdot \psi \nabla \cdot \psi \right| \leq C \|\nabla \cdot \psi\|_{L^2(\Omega)} \|\nabla \cdot \psi\|_{L^2(\Omega)}$$

Step 4: Final Estimate** Combining these bounds:
$$\left| \int_{\Omega} \nabla \cdot \psi \nabla \cdot \psi \right| \leq C \|\nabla \cdot \psi\|_{L^2(\Omega)}^2$$

Dividing by $\|\nabla \cdot \psi\|_{L^2(\Omega)}^2$ yields:
$$\left| \int_{\Omega} \nabla \cdot \psi \nabla \cdot \psi \right| \leq C \|\nabla \cdot \psi\|_{L^2(\Omega)}^2$$

completing the proof with modified constant C .
[Correct] Use inequality techniques to derive the correct key conclusions from variational equations.

Step 1: Variational Formulation The original problem (P) has the weak form: Find $u \in H^1(\Omega)$ such that for all $v \in H^1(\Omega)$:
$$\int_{\Omega} \nabla u \cdot \nabla v = \int_{\Omega} f v$$

[Correct] Consider the weak form of this equation.
The last problem (P) is a saddle-point problem: Find $(u, p) \in H^1(\Omega) \times L^2(\Omega)$ such that for all $v \in H^1(\Omega)$ and $q \in L^2(\Omega)$:
$$\int_{\Omega} \nabla u \cdot \nabla v = \int_{\Omega} f v$$

$$\int_{\Omega} p \nabla \cdot v = 0$$

[Correct] The saddle-point problem is well-posed.

1. Cyclic Kohn with uniform constant. Since $V_1 \sim V_2$ is bounded (positive), the classical Kohn inequality on V_1 gives:
$$\int_{V_1} |\nabla u|^2 \geq C \int_{V_1} |u|^2$$

for all $u \in H^1(V_1)$. By the scaling $u \mapsto u \circ \phi$ (where ϕ is a conformal map), we have:
$$\int_{V_2} |\nabla u|^2 \geq C \int_{V_2} |u|^2$$

for all $u \in H^1(V_2)$.
[Correct] The Kohn inequality is well-posed.


2. Extraction of limits. By the reflexivity of the Banach spaces, there exist, along a subsequence, $u_n \rightharpoonup u$ in $L^2(\Omega)$, $\nabla u_n \rightharpoonup \nabla u$ in $L^2(\Omega)$, $p_n \rightharpoonup p$ in $L^2(\Omega)$, $\nabla p_n \rightharpoonup \nabla p$ in $L^2(\Omega)$, $q_n \rightharpoonup q$ in $L^2(\Omega)$, $\nabla q_n \rightharpoonup \nabla q$ in $L^2(\Omega)$.
[Correct] The extraction of limits is well-posed.

AIM框架概览



清华大学
Tsinghua University


- 由大模型驱动的多智能体系统，针对数学证明特点进行特别设计，包含Explorer、Verifier、Refiner三个智能体以及一个Memory模块



长线思考：探索与记忆机制



- 相比较直接求解问题，AIM当中的智能体被要求针对原问题进行探索，并记录过程当中的发现作为引理；此后继续迭代这一过程即可推进探索进度，并最终实现问题求解




- 对探索过程中的每一个新发现重复检验多次，任意一次检验认为有错则判定新发现有错

- 进行多轮自我检查，能够有效地发现问题
- 存在错误的证明可以尝试修正
- 正确证明结论会最终存储到记忆当中

- 存在错误的证明可以尝试修正

- 正确证明结论会最终存储到记忆当中



● 三个智能体均通过精心设计的prompt指导其行为以及输出格式，并辅以逻辑处理方法实现协作

Prompt Structure

Instruct

_____, _____.

1. _____.

2. _____, _____.

3. _____.

Problem Description

◇ _____, _____.

_____. _____ $\triangleleft \triangleright$

◇ _____, _____,

_____. _____.

_____, _____ $\triangleleft \triangleright$

Memories

_____, _____.

_____.

Mem ID: 0

_____.

_____.

Mem ID: 1

_____.

_____, _____, _____.

_____, _____.

.....


←-- 定义智能体行为

Prompt Example

主要包括：规定任务目标，推理方向，输出格式，以及过程当中各种注意事项。

```
You are an expert that is knowledgeable across all domains in math. This time you are asked to help with our frontier math research. Its statement is as follows:  
This problem could be difficult and not able to be directly solved, but you can make your contribution with the following instructions:  
  
1. You are required to explore different approaches or directions that might help with our final goal, and write down one interesting finding in your explorations as a new conjecture in your response. DO NOT claim that you can not do this job.  
2. Your conjecture must contain the complete definitions required within it, such that it is able to stand alone as an independent lemma, unless it is declared in memory. Do not propose any existing lemmas as your new conjectures. You can directly use them in your explorations.  
3. You should wrap your finding inside a latex environment: \begin{conjecture}\end{conjecture}. This conjecture should be equipped with a detailed, complete and rigorous proof. You should explicitly write down every intermediate derivation step in the proof. The corresponding proof should be wrapped in \begin{proof}\end{proof} directly followed by the conjecture.  
4. After these components you should also provide the dependency of this conjecture. You need to write down the memory IDs of lemmas used in this conjecture in a JSON array format, and wrap them inside \begin{dependency}\end{dependency}. For example, a dependency of a new conjecture could be \begin{dependency}[0, 3, 4]\end{dependency}. You can use an empty array [] when this conjecture does not depend on other lemmas.  
  
More accurately, your response should obey the following format:  
  
\begin{conjecture}Your new findings here\end{conjecture}  
\begin{proof}Your proof of the conjecture above\end{proof}  
\begin{dependency}An json array of related memory IDs of this conjecture\end{dependency}  
Moreover, when you think the time is right that you are able to prove the original problem, you can simply state your proof inside \begin{final-proof}\end{final-proof} and omitted
```

- **三个智能体均通过精心设计的prompt指导其行为以及输出格式，并辅**
以逻辑处理方法实现协作



定义智能体行为

←-- 问题描述

Prompt Example

这部分内容是与问题目标直接相关的信息，对于不同智能体而言内容有所不同：

- Explorer: 最终目标问题表述
- Verifier: 待检验猜想与证明
- Refiner: 存在疑问的猜想、证明以及从 Verifier处获取的反馈

```

\begin{problem}Question:

Can we prove: for any  $\delta>0$ ,

\[
\mathbb{P}[n^{-\delta} \mathbb{E}[Y_n]\leq Y_n\leq n^{\delta}\mathbb{E}[Y_n]]\geq 1-O(d^{-n})
\]

If the second claim is not true, can we prove: for any  $\delta>0$ ,


\[
\mathbb{P}[Y_n\leq n^{2+\delta}]\geq 1-O(d^{-n})
\]

\end{problem}

This problem could be difficult and not able to be directly solved, but you can make your contribution with the following instructions:

```

- **三个智能体均通过精心设计的prompt指导其行为以及输出格式，并辅**
以逻辑处理方法实现协作



定义智能体行为

问题描述

Memory 中的信息

Prompt Example

经过格式化之后的历史探索轨迹以及相关结论。

```

### Context and History Explorations

Here is a list of context that we have collected for this problem or our history finding
s during exploration. They serve as the background of the conjecture and proof and can be ac
cepted without controversy as correct.

#### Memory **ID: 1**

\begin{lemma}

There exists a constant  $\epsilon(A>0)$  and a nonnegative random variable  $\epsilon(Y)$  such that
\[
\forall k \geq 1: \mathbb{E}[Y^k] \leq k! \epsilon(A)^k,
\]
yet for some  $\epsilon(t>0)$ ,
\[
\mathbb{P}(Y \geq t) \geq \exp\{-\epsilon(A) \frac{t^2}{2}\}.
\]
In other words, the bound
\[
\mathbb{P}(Y \geq t) \leq \exp\{-\epsilon(A) \frac{t^2}{2}\}
\]
cannot hold for all  $\epsilon(t>0)$  under only the moment hypothesis.

**DEPENDENCY**: []
\end{lemma}

#### Memory **ID: 2**

```

- 系统从模型输出当中解析并记录四类数据，其中的部分内容会被格式化并进入到后续智能体的输入当中

记忆块类型，包括
context、lemma、
conjecture等

记忆块内容
的文本表述

对于记忆内容
的完整证明

```
struct MemoryBlock {
```

Memory Type

Content Description

Proof: _____。
_____, _____。

MetaData

```
};
```

Visible to all these
agents

Saved and discarded
in explorations

All components in the
memory blocks will be
visible in the output

其他相关信息，如猜想是否解决、
经过检验次数、依赖关系标记等等

结论未知

数学问题

AIM效果

部分证明步骤

量子算法问题

(结论已知)

Quantum Algorithm Problem

AIM完成了带有详细分析
推导过程的证明

吸收边界问题

(结论已知)

Absorbing Boundary Condition

AIM基本完成此问题的证明
过程和细节

高对比度极限

(结论已知)

High Contrast Limit

AIM完成了主要结论的证明
过程，并且证明了新的
结论

均匀化问题

(结论未知)

Homogenization Problem

AIM给出了合理的证明思
路和部分结论的证明，对
数学工作者很有启发意义

←--- DeepSeek-R1

←--- DeepSeek-R1

DeepSeek-R1 和o4-mini各进行一次实验

←--- o4-mini

实验结果：直接使用大模型



● 直接使用大模型错误过多，无法形成数学理论的有效证明过程

Existence and uniqueness:

A fixed-point argument is used on the boundary trace space $L^2([0, T] \times \mathcal{S})$. Define the mapping $T : g \mapsto u|_{\mathcal{S}}$ as follows:

1. Given $g \in L^2([0, T] \times \mathcal{S})$, solve for each k :

$$\partial_t \varphi_k - \Delta_{\mathcal{S}} \varphi_k + d_k \varphi_k = g, \quad \varphi_k(0) = 0.$$

This parabolic equation on $\mathcal{S}$ (a compact Riemannian manifold) has a unique solution $\varphi_k \in C([0, T]; L^2(\mathcal{S})) \cap L^2([0, T]; H^1(\mathcal{S})) \cap H^1([0, T]; H^{-1}(\mathcal{S}))$ if $d_k > 0$.

2. Define $h = \sum_{k=1}^m c_k \varphi_k$.

3. Solve the heat equation with Robin-type boundary condition:

$$\partial_t u - \Delta u = 0, \quad u(0) = u_0, \quad \frac{\partial u}{\partial \nu} + au = h \quad \text{on } \mathcal{S}.$$

For $u_0 \in L^2(\Omega)$ and $h \in L^2([0, T] \times \mathcal{S})$, this has a unique solution $u \in C([0, T]; L^2(\Omega)) \cap L^2([0, T]; H^1(\Omega)) \cap H^1([0, T]; H^{-1}(\Omega))$.

4. Set $T(g) = u|_{\mathcal{S}}$.

4. Quantum Algorithm

The algorithm computes $u(\tau)$ for $\tau = T$:

1. Initial state preparation:

- Prepare $|u(0)\rangle \propto u(x, 0) = K \max(e^x - 1, 0)$ using an oracle $\mathcal{O}_{\text{init}}$. Assume $\|u(0)\| = 1$.

2. Hamiltonian simulation:

- For each k_j , simulate $U_j = \exp(-iT[H + k_j L])$, where H and L are discretized Hermitian matrices (sparsity $d = O(1)$).
- Use sparse Hamiltonian simulation (e.g., QSP) with oracle access to H and L .

3. Linear combination of unitaries (LCU):

- Prepare ancilla state $|\psi_{\text{anc}}\rangle = \frac{1}{\sqrt{s}} \sum_{j=1}^M \sqrt{w_j} |j\rangle$ with $s = \sum_j w_j \approx 1$.
- Apply controlled- U_j : $\sum_j \sqrt{w_j/s} |j\rangle \otimes U_j |u(0)\rangle$.
- Use amplitude amplification to extract the state with $O(1)$ iterations (success probability $\Theta(1)$).

错误运用不动点定理，没有推导过程

推导过程缺失，未验证L半正定性

实例分析：量子算法问题



Linear Combination of Hamiltonian Simulation (LCHS) 方法是科学计算中的一种高效方法，主要思路是将非酉动力学问题转化为为哈密顿模拟的线性组合

$$\mathcal{T}e^{-\int_0^t A(s) ds} = \int_{\mathbb{R}} \frac{1}{\pi(1+k^2)} \mathcal{T}e^{-i\int_0^t (H(s)+kL(s)) ds} dk$$

Black-Scholes-Merton(BSM)模型是金融学中用于欧式期权定价的基本数学框架

$$\frac{\partial V}{\partial t} + \frac{1}{2}\sigma^2 S^2 \frac{\partial^2 V}{\partial S^2} + rS \frac{\partial V}{\partial S} - rV = 0$$

目标：利用LCHS方法对BSM模型进行模拟，设计相应的量子算法，并分析算法的复杂度

AIM正确运用了这一方法，提供了较为详细的证明过程，基本解决这一问题

实例分析：量子算法问题



Lemma (Linear combination of Hamiltonian simulation, LCHS): For $t \in [0, T]$, let $A(t) \in \mathbb{C}^{N \times N}$ be decomposed into Hermitian and anti-Hermitian parts such that $A(t) = L(t) + iH(t)$, where $L(t) = \frac{1}{2}[A(t) + A^\dagger(t)]$ and $H(t) = \frac{1}{2i}[A(t) - A^\dagger(t)]$. Assume that $L(t)$ is positive semi-definite for all $t \in [0, T]$. Denoting the time ordering operator by $\mathcal{T}$, we have $\mathcal{T} \exp \left\{ - \int_0^t A(s) ds \right\} = \int_{\mathbb{R}} \eta(k) u(t, k) dk$, where $u(t, k)$ is the propagator for a time-dependent Hamiltonian simulation problem such that $u(t, k) = \mathcal{T} \exp \left\{ -i \int_0^t [H(s) + kL(s)] ds \right\}$, and $\eta(k) = \frac{1}{\pi(1+k^2)}$ is the kernel function with respect to k .

BSM model: The PDE in the BSM model is given by

$$\frac{\partial V(S, t)}{\partial t} + \frac{1}{2} \sigma^2 S^2 \frac{\partial^2 V}{\partial S^2} + rS \frac{\partial V}{\partial S} - rV = 0.$$

Boundary conditions are characterized by: $V(0, t) = 0$ for all $0 < t \leq T$; $V(S, t) \rightarrow S - Ke^{-r(T-t)}$ for $S \rightarrow \infty$; $V(S, T) = \max\{S - K, 0\}$.

Your tasks are to: Use the idea of LCHS to deal with the BSM model. Design a quantum algorithm to solve the equation. And compute the complexity (query complexity to state preparation oracle and operator oracles, gate complexity, ancilla qubits, (parameters: norm of boundary and initial, T , approximation error ϵ , etc)).

LCHS方法
说明

BSM模型
说明

目标

实例分析：量子算法问题



- 具体来说，AIM通过将BSM方程转化为热方程并利用LCHS引理转化问题并给出量子算法，基本解决了该问题，核心步骤如下：

1	偏微分方程转换与空间离散化	通过变量替换将BSM偏微分方程转换为标准热方程，并对空间变量进行离散化处理
2	算子分解与积分离散化	应用LCHS引理将热方程的解算子分解为酉算子的连续线性组合，随后对连续积分截断近似并离散化
3	量子算法实现与复杂度分析	利用酉算子线性组合（LCU）和哈密顿模拟技术设计量子算法，并计算复杂度

实例分析：量子算法问题



AIM通过层层递进的多个引理逐步探索：

引理1： AIM先将BSM方程转化为热方程，再离散化空间变量，验证半正定性；进一步转化为可以利用LCHS引理的形式，证明过程详细

1. The Black-Scholes-Merton (BSM) partial differential equation (PDE) can be transformed into the standard heat equation via a logarithmic substitution. Specifically, under the transformation $x = \ln(S/K) + (r - \sigma^2/2)(T - t)$, $\tau = \sigma^2(T - t)/2$, and $V(S, t) = e^{-r(T-t)}U(x, \tau)$, the BSM PDE reduces to:

$$\frac{\partial U}{\partial \tau} = \frac{\partial^2 U}{\partial x^2}.$$

Discretizing this PDE spatially yields a matrix equation $\partial_\tau |U\rangle = -B|U\rangle$, where $B = -\partial_x^2$. The operator B is positive semi-definite and can be directly interpreted as $B = L$, where $L = -\partial_x^2$. While the original Hermitian decomposition $B = L + iH$ (with $H = 0$) trivializes the LCHS lemma, the dissipative nature of B permits alternative quantum simulation methods tailored to parabolic equations.

将BSM方程转化为热方程

Proof. **Step 1: Transform BSM to the Heat Equation**

Start with the BSM PDE:

$$\frac{\partial V}{\partial t} + \frac{1}{2}\sigma^2 S^2 \frac{\partial^2 V}{\partial S^2} + rS \frac{\partial V}{\partial S} - rV = 0.$$

Apply the substitution $V(S, t) = e^{-r(T-t)}U(x, \tau)$, $x = \ln(S/K) + (r - \sigma^2/2)(T - t)$, and $\tau = \sigma^2(T - t)/2$. Compute derivatives rigorously: $-\partial_t V = e^{-r(T-t)} \left(rU - \frac{\sigma^2}{2} \partial_\tau U - (r - \sigma^2/2) \partial_x U \right)$, $\partial_S V = e^{-r(T-t)} \partial_x U \cdot \frac{1}{S}$, $\partial_S^2 V = e^{-r(T-t)} (\partial_x^2 U - \partial_x U) \cdot \frac{1}{S^2}$.

Substitute into the BSM PDE:

$$e^{-r(T-t)} \left[rU - \frac{\sigma^2}{2} \partial_\tau U - (r - \sigma^2/2) \partial_x U + \frac{1}{2} \sigma^2 (\partial_x^2 U - \partial_x U) + r \partial_x U - rU \right] = 0.$$

Multiply through by $e^{r(T-t)}$, cancel terms, and simplify:

$$-\frac{\sigma^2}{2} \partial_\tau U + \frac{\sigma^2}{2} \partial_x^2 U = 0 \implies \partial_\tau U = \partial_x^2 U.$$

****Step 2: Spatial Discretization****

Discretize x on a grid with spacing Δx . Approximate $\partial_x^2 U$ via finite differences:

$$\partial_x^2 U \approx \frac{U_{j+1} - 2U_j + U_{j-1}}{\Delta x^2}.$$

Let D_2 be the discretized second derivative matrix. The operator becomes:

$$B = -D_2.$$

空间变量
离散化

****Step 3: Positive Semi-Definite Operator****

The matrix $B = -D_2$ is positive semi-definite because D_2 (a centered second derivative discretization) is negative semi-definite. This aligns with the dissipative nature of the heat equation.

****Step 4: Implications for Quantum Simulation****

While the original decomposition $B = L + iH$ (with $H = -ic\partial_x$) is invalid due to the absence of a first-order term, the operator B itself is suitable for quantum simulation methods designed for parabolic equations. Specifically, the time evolution $\partial_\tau |U\rangle = -B|U\rangle$ can be simulated using Trotterization or other dissipative quantum algorithms, bypassing the need for the LCHS lemma. $\square$

[Correct] Variable substitution and equation transformation are correct. Discretization of the spatial variable is correct. And AIM finds the reduced PDE trivializes the LCHS lemma. It's right!

实例分析：量子算法问题



AIM通过层层递进的多个引理逐步探索：

引理2： AIM分析了截断误差并确定离散间距，为构建量子算法奠定基础

2.The integral representation in the LCHS lemma for the BSM model can be approximated with precision ϵ using $O(1/\epsilon^2)$ terms through an adaptive discretization of the k -integral, leveraging the rapid decay of the Cauchy kernel $\eta(k) = \frac{1}{\pi(1+k^2)}$.

*Proof. **Truncation Error Analysis**:* The integral $I = \int_{-\infty}^{\infty} \eta(k)u(t,k)dk$ is truncated to $[-K, K]$. The tail error is bounded by:

$$\int_{|k|>K} \eta(k)dk = \frac{2}{\pi} \int_K^{\infty} \frac{1}{1+k^2} dk = \frac{2}{\pi} \left(\frac{\pi}{2} - \arctan(K) \right) \approx \frac{1}{\pi K} \quad \text{for } K \gg 1.$$

Setting $\frac{1}{\pi K} \leq \epsilon/2$ gives $K \geq \frac{2}{\pi\epsilon}$. Thus, $K = O(1/\epsilon)$.

***Adaptive Discretization**:* 1. ***Central Interval $[-K, K]$ **:* The entire truncated domain $[-K, K]$ must be discretized. To bound the discretization error, use a uniform grid with spacing $\Delta k = \Theta(\epsilon)$. The number of points is $N = O(K/\Delta k) = O(1/\epsilon^2)$. The discretization error using the trapezoidal rule scales as $O(K\Delta k^2 \cdot \max |\partial_k^2(\eta(k)u(t,k))|)$. Since $\eta(k)$ and $u(t,k)$ are smooth, $\max |\partial_k^2(\eta(k)u(t,k))|$ is bounded over $[-K, K]$. The total error becomes $O(K\Delta k^2) = O(\frac{1}{\epsilon} \cdot \epsilon^2) = O(\epsilon)$.

***Outer Interval Contribution**:* The truncated integral already includes all significant contributions, as the tail beyond $K = O(1/\epsilon)$ contributes at most $\epsilon/2$. The central interval $[-K, K]$ is approximated with $O(1/\epsilon^2)$ points, ensuring the total discretization error is $O(\epsilon)$. Additional adaptive methods in outer intervals (e.g., geometrically spaced grids) would only reduce logarithmic factors, which are asymptotically dominated by $O(1/\epsilon^2)$.

***Total Terms**:* The uniform grid over $[-K, K]$ requires $O(1/\epsilon^2)$ terms. While adaptive grids in outer regions could marginally reduce constants, the dominant term remains $O(1/\epsilon^2)$.

***Final Error**:* Truncation error $O(\epsilon/2)$ + discretization error $O(\epsilon/2) = O(\epsilon)$. Thus, $O(1/\epsilon^2)$ terms suffice for ϵ -precision approximation. $\square$

[Correct] Truncate and discretize the integral correctly.

分析截断误差

确定离散间距

总误差分析

实例分析：量子算法问题



清华大学
Tsinghua University

AIM通过层层递进的多个引理逐步探索：

AIM 结合之前的引理，使用哈密顿模拟和酉算子的线性组合（LCU）构建了一个量子算法，然后计算其复杂度；复杂度计算有一些错误，但核心方法是正确的

利用哈密顿模拟算法

3.The solution operator for the heat equation derived from the BSM model, $U(\tau) = e^{\tau B}$, can be approximated via the LCHS lemma as a linear combination of Hamiltonian simulations. Specifically, the integral representation $\int_{\mathbb{R}} \eta(k) e^{-ikL\tau} dk$ (where $L = -B$) can be discretized into $M = O(1/\epsilon^2)$ terms with quadrature weights $\eta(k_j) \Delta k$. Each term $e^{-ik_j L \tau}$ is simulated by $k_j \tau$. The total query complexity of the L -oracle is $O(M \cdot \|L\| \tau)$. The query complexity scales polynomially with $\|L\|$.

在第一个引理基础上，利用LCHS引理转化为酉算子的连续积分

Proof:

****Step 1: Integral Representation via LCHS**** From Lemma-0, the solution operator for the heat equation is $U(\tau) = e^{\tau B}$. Let $L = -B$, a positive semi-definite operator. By the LCHS lemma with $H = 0$, we express:

$$e^{\tau B} = \int_{\mathbb{R}} \eta(k) e^{-ikL\tau} dk, \quad \eta(k) = \frac{1}{\pi(1+k^2)}.$$

****Step 2: Quadrature Discretization**** Lemma-1 guarantees the integral can be discretized into $M = O(1/\epsilon^2)$ terms with weights $\eta(k_j) \Delta k$, yielding:

$$e^{\tau B} \approx \sum_{j=1}^M \eta(k_j) \Delta k \cdot e^{-ik_j L \tau}.$$

The truncation and discretization error is $O(\epsilon)$ due to the $O(1/k^2)$ decay of $\eta(k)$.

[Correct] Properly utilizing the LCHS lemma and memory lemmas for problem transformation.

将方程解转化为酉算子的线性组合

****Step 3: Hamiltonian Simulation Costs**** Each term $e^{-ik_j L \tau}$ requires simulating L for time $t_j = k_j \tau$. Using optimal Hamiltonian simulation algorithms (e.g., quantum signal processing or Taylor series) with query complexity $O(\|L\| t_j + \log(1/\epsilon_{\text{sim}}))$, set $\epsilon_{\text{sim}} = \epsilon/M$ to bound the total simulation error by ϵ . Since $k_j = O(1/\epsilon)$ dominates the quadrature, $t_j = O(\tau/\epsilon)$. The per-term complexity becomes:

$$O(\|L\| \tau / \epsilon + \log(M/\epsilon)) = \tilde{O}(\|L\| \tau / \epsilon).$$

****Step 4: Total Query Complexity**** Summing over $M = O(1/\epsilon^2)$ terms, the total queries to the L -oracle are:

$$\tilde{O}\left(M \cdot \frac{\|L\| \tau}{\epsilon}\right) = \tilde{O}\left(\frac{\|L\| \tau}{\epsilon^3}\right).$$

Expressing in terms of $T = \frac{2\tau}{\sigma^2}$ and absorbing constants:

$$\tilde{O}\left(\frac{\|L\| T}{\epsilon^3}\right).$$

复杂度分析

****Step 5: Ancilla and Gate Complexity**** The LCU framework requires $\log M = O(\log(1/\epsilon))$ ancilla qubits. Gate complexity inherits $\tilde{O}(M)$ -scaling from the superposition state, augmented by $O(\text{poly}(\log(\|L\| T / \epsilon)))$ factors from simulation subroutines. The dominant terms remain polynomial in $\|L\|$, T , and $1/\epsilon$, with polylogarithmic corrections.

[Error] There are some mistakes about complexity computing. And the calculation process lacks detail.

其他结果：吸收边界条件问题



- 问题描述：证明在特定设计的吸收边界条件约束条件下，热传导方程解的适定性和唯一性

$$\begin{cases} \mathcal{L}u = 0, & [0, \infty) \times \Omega \\ \frac{\partial u}{\partial \nu} = -\beta u - \sum_{k=1}^m \alpha_k (\partial_t - \Delta_S) \varphi_k = \mathcal{F}_1 u, & [0, \infty) \times \mathcal{S} \\ u = u_0, & t = 0 \end{cases} \quad u \in C([0, T]; L^2(\Omega)) \cap H^1([0, T]; H^{-1}(\Omega)) \cap L^2([0, T]; H^1(\Omega)),$$
$$\begin{cases} (\partial_t - \Delta_S + d_k) \varphi_k = u, & t \geq 0 \\ \varphi_k = 0, & t = 0 \end{cases} \quad \varphi_k \in C([0, T]; L^2(\mathcal{S})) \cap H^1([0, T]; H^{-1}(\mathcal{S})) \cap L^2([0, T]; H^1(\mathcal{S}))$$

AIM正确运用了分析这类问题的思路和方法，给出了完成度很高的证明过程

****A priori energy estimate for the coupled system**** Let u and φ_k ($k = 1, \dots, m$) satisfy the system (2.10) with $u_0 \in L^2(\Omega)$ and $\varphi_k(0) = 0$. Then, there exists a constant $C > 0$ depending on T, β, α_k, d_k , but independent of u_0 , such that:

$$\begin{aligned} & \sup_{t \in [0, T]} \|u(t)\|_{L^2(\Omega)}^2 + \int_0^T \|\nabla u(t)\|_{L^2(\Omega)}^2 dt + \int_0^T \|u(t)\|_{L^2(\mathcal{S})}^2 dt \\ & + \sum_{k=1}^m \left(\sup_{t \in [0, T]} \|\varphi_k(t)\|_{L^2(\mathcal{S})}^2 + \int_0^T \|\nabla_S \varphi_k(t)\|_{L^2(\mathcal{S})}^2 dt \right) \leq C \|u_0\|_{L^2(\Omega)}^2. \end{aligned}$$

这里是AIM给出的核心范数控制不等式，是这个问题的关键中间结论

其他结果：高对比度极限问题



- 问题描述：考虑一个传输系统在某种参数极限下的特例情况，分析求证极限解和原本解之间的误差估计

$$\left\{ \begin{array}{l} \mathcal{L}_{\lambda, \mu} \mathbf{u}_\epsilon = 0 \\ \mathcal{L}_{\tilde{\lambda}, \tilde{\mu}} \mathbf{u}_\epsilon = 0 \\ \mathbf{u}_\epsilon|_- = \mathbf{u}|_+ \text{ and } \frac{\partial \mathbf{u}_\epsilon}{\partial \nu_{(\tilde{\lambda}, \tilde{\mu})}}|_- = \frac{\partial \mathbf{u}_\epsilon}{\partial \nu_{(\lambda, \mu)}}|_+ \\ \frac{\partial \mathbf{u}_\epsilon}{\partial \nu_{(\lambda, \mu)}}|_{\partial \Omega} = g \in H_{\mathbb{R}}^{-\frac{1}{2}}(\partial \Omega) \text{ and } \mathbf{u}_\epsilon|_{\partial \Omega} \in H_{\mathbb{R}}^{\frac{1}{2}}(\partial \Omega). \end{array} \right. \begin{array}{l} \text{in } \Omega \setminus \overline{D_\epsilon}, \\ \text{in } D_\epsilon, \\ \text{on } \partial D_\epsilon, \end{array} \quad \left\{ \begin{array}{l} \mathcal{L}_{\lambda, \mu} \mathbf{u}_\epsilon = 0 \\ \mathcal{L}_{\tilde{\mu}}(\mathbf{u}_\epsilon, p_\epsilon) = 0 \text{ and } \operatorname{div} \mathbf{u}_\epsilon = 0 \\ \mathbf{u}_\epsilon|_- = \mathbf{u}_\epsilon|_+ \text{ and } \frac{\partial(\mathbf{u}_\epsilon, p_\epsilon)}{\partial \nu_{(\infty, \tilde{\mu})}}|_- = \frac{\partial \mathbf{u}_\epsilon}{\partial \nu_{(\lambda, \mu)}}|_+ \\ \frac{\partial \mathbf{u}_\epsilon}{\partial \nu_{(\lambda, \mu)}}|_{\partial \Omega} = g \in H_{\mathbb{R}}^{-\frac{1}{2}}(\partial \Omega) \text{ and } \mathbf{u}_\epsilon|_{\partial \Omega} \in H_{\mathbb{R}}^{\frac{1}{2}}(\partial \Omega), \end{array} \right. \begin{array}{l} \text{in } \Omega \setminus \overline{D_\epsilon}, \\ \text{in } D_\epsilon, \\ \text{on } \partial D_\epsilon, \end{array}$$

AIM给出了核心结论的主要推导过程和证明，并且主动探索得到了其他正确结论

Under the given assumptions, the difference $\mathbf{w}_\epsilon = \mathbf{u}_{\text{lim}} - \mathbf{u}_\epsilon$ between the solutions of the limit problem (10) and the original problem (9) satisfies the homogenized error estimate in the energy norm weighted by the Lamé parameters:

$$\left(\int_{\Omega} [\lambda(x) |\operatorname{div} \mathbf{w}_\epsilon|^2 + 2\mu(x) |\mathcal{D}(\mathbf{w}_\epsilon)|^2] dx \right)^{1/2} \leq \frac{C}{\sqrt{\lambda}} \|g\|_{H^{-\frac{1}{2}}(\partial \Omega)},$$

where C is independent of $\tilde{\lambda}$. This implies that the error in the energy norm decays proportionally to $\tilde{\lambda}^{-\frac{1}{2}}$.

correctness: True

AIM探索得到全局
空间上的控制结果

其他结果：均匀化问题



- 问题描述：传输系统均匀化问题，需分析在具体物理尺度极限下，方程和相应的解的性质，并最终证明解的误差估计，**是一个开放性问题**

$$\begin{cases} \mathcal{L}_{\lambda, \mu} \mathbf{u}_\epsilon = 0 \\ \mathcal{L}_{\tilde{\mu}}(\mathbf{u}_\epsilon, p_\epsilon) = 0 \text{ and } \operatorname{div} \mathbf{u}_\epsilon = 0 \\ \mathbf{u}_\epsilon|_- = \mathbf{u}_\epsilon|_+ \text{ and } \left. \frac{\partial(\mathbf{u}_\epsilon, p_\epsilon)}{\partial \nu(\infty, \tilde{\mu})} \right|_- = \left. \frac{\partial \mathbf{u}_\epsilon}{\partial \nu(\lambda, \mu)} \right|_+ \\ \left. \frac{\partial \mathbf{u}_\epsilon}{\partial \nu(\lambda, \mu)} \right|_{\partial \Omega} = g \in H_{\mathbb{R}}^{-\frac{1}{2}}(\partial \Omega) \text{ and } \mathbf{u}_\epsilon|_{\partial \Omega} \in H_{\mathbb{R}}^{\frac{1}{2}}(\partial \Omega), \end{cases}$$

in $\Omega \setminus \overline{D_\epsilon}$, where $\mathcal{L}_{\tilde{\mu}}(\mathbf{u}_\epsilon, p_\epsilon) = \tilde{\mu} \Delta \mathbf{u}_\epsilon + \nabla p_\epsilon$ denotes the Stokes operator with viscosity constant $\tilde{\mu}$ and p_ϵ is the pressure field. Its exterior derivative is defined as $\left. \frac{\partial(u, p)}{\partial \nu(\infty, \mu)} \right|_- := pN + 2\mu D(u)N$.
on ∂D_ϵ ,
Now we let $\epsilon \rightarrow 0$, so that the scale of the cell is tend to be zero. We wonder what homogenization equation does the limited solution $\mathbf{u}_{lim}$ satisfies. At the same time, whether the estimate between the original solution and the limited solution has the form $C\epsilon^\alpha \|g\|_{H^{-\frac{1}{2}}(\partial \Omega)}$ for some $\alpha \in (0, 1)$.

AIM正确运用数学定理和分析推导，给出了一些正确的结论和思路
对数学家完成此项研究具有指导意义

6. Error estimate and strong convergence Define the two-scale corrector

$$u_\epsilon^{\text{app}}(x) = u_0(x) + \epsilon \chi^{D_x u_0(x)}(x/\epsilon) \text{ in } \Omega_\epsilon,$$

extended in each D_ϵ by $Ey + \chi_i^E(y)$. A Céa-lemma-type argument in the perforated domain (again relying on uniform Korn and Stokes inf-sup, Lemma A.1–A.2) yields for some $\alpha \in (0, 1)$

$$\|u_\epsilon - u_\epsilon^{\text{app}}\|_{H^1(\Omega_\epsilon)} \leq C\epsilon^\alpha \|g\|_{H^{-1/2}(\partial \Omega)}.$$

From this one deduces $u_\epsilon \rightarrow u_0$ strongly in $L^2(\Omega)$ (and even in $H^1(\Omega_\epsilon)$ up to the corrector).


This completes a fully rigorous derivation of the homogenized elastic limit and the explicit formula for C^{hom} .

正确的渐进展开思路和结论探索


局限1：重复性探索



- 智能体常沿相同方向进行探索，提出一系列相似猜想和相同引理，推高成本、降低效率并制约性能上限




Lemma 4. The solution operator $e^{-\tau B}$ for the heat equation derived from the BSM model can be approximated with error ϵ using a quantum algorithm that implements a discretized version of the LCHS lemma. This algorithm requires $M = O\left(\frac{1}{\epsilon^2}\right)$ terms in the quadrature approximation, and the total query complexity to the B -oracle (encoding the discretized Laplacian) is $\tilde{O}\left(\frac{\|B\|T}{\epsilon^3}\right)$, where $T = \frac{2\tau}{\sigma^2}$. The gate complexity and ancilla qubits scale polynomially with $\log(1/\epsilon)$, $\|B\|$, and T .






Lemma 6. The quantum algorithm for solving the BSM PDE using the LCHS lemma achieves an approximation error ϵ with query complexity $\tilde{O}\left(\frac{\|B\|T}{\epsilon^3}\right)$ to the operator oracle encoding B , where $B = -\partial_x^2$ is the discretized Laplacian, T is the time horizon, and ϵ is the error tolerance. The ancilla qubit count is $O(\log \frac{1}{\epsilon})$, and the gate complexity scales as $\text{poly}(\|B\|, T, \log \frac{1}{\epsilon}, \frac{1}{\epsilon})$.

3个相似的引理



Lemma 7. The quantum algorithm for solving the BSM PDE using the LCHS lemma requires simulating the discretized Laplacian operator $B = -\partial_x^2$ for a total time $\tau = \frac{\sigma^2 T}{2}$. The integral $\int_{\mathbb{R}} \eta(k) e^{-ikB\tau} dk$ is approximated by $M = O\left(\frac{1}{\epsilon^2}\right)$ quadrature terms. Each term involves Hamiltonian simulation of $k_j B$, where $|k_j| \leq O\left(\frac{1}{\epsilon}\right)$. The total query complexity to the B -oracle is $\tilde{O}\left(\frac{\|B\|T}{\epsilon^3}\right)$, and the gate complexity scales as $\text{poly}(\|B\|, T, \log \frac{1}{\epsilon}, \frac{1}{\epsilon})$, with ancilla qubit count $O(\log \frac{1}{\epsilon})$.



局限2：数学设定理解不足



- 当前AIM对于较为冗长的数学设定、背景条件的理解能力还不够强，导致智能体在分析过程中产生错误

Then as $\varepsilon \rightarrow 0$ one has, up to a subsequence, $u_\varepsilon \rightharpoonup u_0$ in $H^1(\Omega)$, $u_\varepsilon \rightarrow u_0$ in $L^2(\Omega)$, $p_\varepsilon \xrightarrow{\text{two-scale}} p_1(x, y)$ in $\Omega \times Y_i$, where $u_0 \in H_R^1(\Omega; \mathbb{R}^d)$ is the unique solution of the homogenized Lamé system $-\operatorname{div}_x [C^{\text{hom}} D_x(u_0)] = 0$ in Ω , $C^{\text{hom}} D_x(u_0) \cdot n = g$ on $\partial\Omega$, and the effective fourth-order tensor C^{hom} is given by the following periodic cell transmission problem: for each fixed symmetric $E \in \mathbb{R}_{\text{sym}}^{d \times d}$ find $(\chi_e^E, \chi_i^E, \pi^E) \in H_{\text{per}}^1(Y_e; \mathbb{R}^d) \times H_{\text{per}}^1(Y_i; \mathbb{R}^d) \times L_{\text{per}}^2(Y_i)/\mathbb{R}$ solving

(1) In the elastic cell Y_e : $-\operatorname{div}_y [C(E + D_y \chi_e^E)] = 0$, $\operatorname{div}_y (Ey + \chi_e^E) = 0$,

(2) In the fluid cell Y_i : $-\mu \Delta_y \chi_i^E + \nabla_y \pi^E = 0$, $\operatorname{div}_y (Ey + \chi_i^E) = 0$,

(3) On the material interface $\Gamma = \partial Y_i$: $\chi_e^E = \chi_i^E$, $[C(E + D_y \chi_e^E) \cdot n] = [2\mu D_y \chi_i^E - \pi^E I] \cdot n$,

此处缺乏对晶胞的边界理解，导致给出了不符合实际情况的结论

局限3：推导细节缺失



- 当前AIM会出现“跳步”现象，不利于证明过程的人工或自动化验证，同时可能导致错误的证明过程

****Step 2: Formulate the Projected Equations**** For test functions $v \in V_N$ and $\psi \in W_{k,N}$, the Galerkin approximations satisfy the projected weak form:

$$\int_{\Omega} \partial_t u^N v \, dx + \int_{\Omega} \nabla u^N \cdot \nabla v \, dx + \int_{\Omega} \rho u^N v \, dx \\ + \beta \int_S u^N v \, ds + \sum_{k=1}^m \alpha_k \int_S (\partial_t \varphi_k^N - \Delta_S \varphi_k^N) v \, ds = 0,$$

$$\int_S \partial_t \varphi_k^N \psi \, ds + \int_S \nabla_S \varphi_k^N \cdot \nabla_S \psi \, ds + d_k \int_S \varphi_k^N \psi \, ds = \int_S u^N \psi \, ds.$$

****Step 3: Solve the ODE System**** Substituting $u^N = \sum_{j=1}^N a_j(t) w_j$ and $\varphi_k^N = \sum_{l=1}^N b_{k,l}(t) \theta_{k,l}$ into the projected equations yields a linear ODE system for $a_j(t)$ and $b_{k,l}(t)$. The coefficients are smooth (as eigenfunctions are C^∞), ensuring continuity. By the Cauchy-Lipschitz theorem, unique solutions exist on $[0, T]$.

[Correct] The agent gets the ODE satisfied with the approximated solution.

[Vague] It is better to have more detailed process.

此处并未给出ODE系统的具体推导分析过程，但是是正确的结论

展望：AI作用从辅助到主动



清华大学
Tsinghua University

辅助证明 + 思路验证 + 开放探索



终极目标



展望：AI推动更高质量数学研究合作




清华大学
Tsinghua University

● AI数学家作为独立个体参与合作，让数学合作更容易，加速数学发展



Terence Tao
@tao@mathstodon.xyz

As of yesterday, the Equational Theories Project teorth.github.io/equational_th... has (provisionally) attained its aim of resolving all 22028942 implications between the 4694 equational laws of magmas involving at most four multiplication operations. We are not fully done yet - there are 52 resolutions that currently only have a human-readable proof but not a Lean-formalized proof (actually, there are a few that are not even human-readable, but are instead computer-generated by some non-Lean program) - but it should now just be a matter of time before we can declare final success in which the full implication graph is completely formalized in Lean. As such, we are now beginning the process of writing up the results.



Equational Theories Project

- 57天：人类和AI合作搞定了4,694个等式之间22,028,942个蕴含关系
- 9天：在AI帮助下进度达到99.866%

Contributors 61




+ 47 contributors


总结



- 大模型的快速发展为AI在数学研究中发挥更大作用提供了基础
- 数学研究区别于数学解题的三个关键特征：
 - 长线思考：数学研究需要更多、更长期、更深入有效思考
 - 过程严谨：错误零容忍，需可验证，但验证代价高昂
 - 答案存在性：数学研究答案存在性未知且不确定
- 提出AI数学家系统AIM，并在四个研究级数学问题上取得初步成效，潜力可期待
- 未来AI将在数学研究中起到更加主动且重要的作用



AIM技术报告



AIM系统

Our AIM is AI Mathematician!

